Hepatitis C Could I be at risk?



Contents

Page no

What is hepatitis C?	3
How could I get hepatitis C?	3
What are the symptoms?	5
How might the virus affect me in the long term?	6
How do I know if I have the virus?	6
Where can I get a test?	7
What if the test result is negative?	7
What if the test result is positive?	7
Is treatment available?	8
How can I help myself if I have the virus?	8
How can I avoid becoming infected or passing on the virus to others?	8
Information and support	10

What is hepatitis C?

Hepatitis C is a virus that can damage the liver and can lead to years of ill health. Unlike hepatitis A and B, there is no vaccine to protect against the hepatitis C virus (HCV).

How could I get hepatitis C?

Hepatitis C is carried in the blood and can be found in some other body fluids. But blood (even in tiny or invisible amounts) is the main source of infection. The virus is spread mainly by the blood of an infected person entering another person's bloodstream. You can't catch it through everyday contact such as holding hands, hugging or kissing, or through sharing toilets, crockery or kitchen utensils.

Hepatitis C is passed on by:

 sharing equipment for injecting drugs, even if you did this only once or twice, or a long time ago. Needles and syringes are the greatest risk, but spoons, filters or straws for inhaling cocaine could also carry infection, as could water; receiving a blood transfusion (before September 1991), or blood products like clotting factors used to treat haemophilia (before 1986). All blood used in Northern Ireland is now screened for hepatitis C.

Less commonly, hepatitis C can be passed on by:

- a hepatitis C positive mother to her baby, before or during birth;
- unprotected sex (without a condom) with someone who has the virus;
- living in countries where hepatitis C is more common and there is an increased risk of it being passed on through medical and dental procedures and where equipment may not be sterilised properly;
- tattooing, ear piercing, body piercing or acupuncture with equipment that is not sterile;
- sharing razors or toothbrushes contaminated with blood from someone who has the virus.

Will I have symptoms?

Many people who have the hepatitis C virus have no signs or symptoms at all – for years or even decades. But even without symptoms you can still pass the virus on to others.

How do I know if I have the virus?

If you think you could have been in contact with the hepatitis C virus at any time in the past, you can have a simple blood test to find out if you may have been infected.

Where can I get a test?

Your GP will be able to carry out tests for hepatitis C. Sexual health/Genito Urinary Medicine (GUM) clinics may also offer testing as may some drug treatment services. (See page 11 of Information and support for details on how to contact these services.)

What if the test result is negative?

This probably means that you have never been in contact with the hepatitis C virus. However, it can take up to six months for your body to react to the virus, so the test result may be negative if you have been infected recently. Your GP may advise you to have a second test.

What if the test result is positive?

A blood test will tell you whether you have been infected with the virus at any stage in the past and whether you are still carrying the virus. You can be referred to a specialist to see how it is affecting your liver.

How might the virus affect me in the long term?

Around one in five people who become infected with hepatitis C will recover completely without treatment. However, most people who become infected remain so for a very long time. This may affect them in different ways:

some people can remain well throughout their life;

- some people experience a range of symptoms;
- about one in five people will develop severe liver damage (cirrhosis) after about 20 years;
- in some people, cirrhosis can lead to liver cancer or complete liver failure.

Is treatment available?

Treatment for hepatitis C is available and is improving all the time. Drug therapy is available that can cure the infection in about half of the people treated. Your GP and specialist will discuss which treatment is most appropriate for you.

How can I help myself if I have the virus?

If you drink alcohol, the most important thing you can do to minimise damage to your liver is to limit the amount of alcohol you drink or, better still, to cut out alcohol altogether.

How can I avoid becoming infected or passing on the virus to others?

- Avoid any direct contact with other people's blood.
- Avoid sharing any toiletry equipment that could have been contaminated with blood, eg razors, toothbrushes, nail scissors, tweezers.
- If you or your sexual partner has hepatitis C, there is a small risk that it could be passed on during sex. Condoms minimise the risk. Using condoms will also help to protect you against unwanted pregnancy, HIV and many other sexually transmitted infections.
- If you are considering having ear piercing, body piercing, a tattoo or acupuncture, check that equipment such as needles and ink pots are not being re-used and are sterile.
- If you inject drugs, NEVER share or pass on drug injecting equipment. This includes needles, syringes, spoons, water, filters or anything else that might be contaminated with blood. Contact the National Drugs Helpline (FRANK) on 0800 77 66 00 for details of drug services in your area and safer injecting advice.

Information and support

If you have any questions, you can talk to your GP or nurse.

- Sexual health/Genito Urinary Medicine (GUM) clinics provide free, confidential advice. Please phone for further information.
 Royal Victoria Hospital, Belfast: Tel: 028 9063 4050/028 9063 4054
 Causeway Hospital, Coleraine: Tel: 028 7034 6028
 Daisy Hill Hospital, Newry: Tel: 028 3083 5050 (calls returned within 2 working days)
 Altnagelvin Hospital, Londonderry: Tel: 028 7161 1269
 Downe Hospital, Downpatrick: Tel: 028 4483 8133
- **www.drugsalcohol.info** lists the drug and alcohol treatment services throughout Northern Ireland. Some of these may offer testing but you may need to be referred by your GP.
- Hepatitis C information line is for anyone with concerns about hepatitis C.
 Helpline: 0800 451 451 (8am–8pm, Mon–Fri; 10am–4pm, Sat–Sun)
 Textphone: 0800 085 0859

- British Liver Trust (BLT) provides information on all forms of liver disease.
 Helpline: 0800 652 7330 (9am–5pm, Mon–Fri) www.britishlivertrust.org.uk

Tel: 0845 223 4424 (10.30am–4.30pm, Mon–Fri) www.hepctrust.org.uk

- UK Hepatitis C Resource Centre provides information for people with hepatitis C. Tel: 0870 242 2467 (9am–5pm, Mon–Fri) www.hepccentre.org.uk
- **The Haemophilia Society** provides information and support for people living with or affected by haemophilia who are concerned about hepatitis C.

Helpline: 0800 018 6068 (10am–3pm, Tues–Fri, all calls returned within 5 working days) www.haemophilia.org.uk

 Children's Liver Disease Foundation (CLDF) provides support for children with liver disease. Tel: 0121 212 3839 www.childliverdisease.org • **RVH Liver Support Group** offers support to those in Northern Ireland living with liver disease and is an independent group affiliated to the British Liver Trust.

Tel: 077 3771 8493 www.rvhliversupportgroup.org

- Northern Ireland Hepatitis C Managed Clinical Network provides information on hepatitis C. www.hepcni.net
- National Drugs Helpline (FRANK) provides information and advice for drug users and anyone concerned about drugs.

Helpline: 0800 77 66 00 (24 hours a day, 365 days a year) Textphone: 0800 917 8765 www.talktofrank.com

For more information, visit the Public Health Agency website, www.publichealth.hscni.net or the Department of Health, Social Services and Public Safety website, www.dhsspsni.gov.uk/phealth or the NHS hepatitis C website, www.hepc.nhs.uk

This leaflet is also available as PDF files in other languages from: www.publichealth.hscni.net www.dhsspsni.gov.uk/phealth







Published by the Public Health Agency, Ormeau Avenue Unit, 18 Ormeau Avenue, Belfast BT2 8HS. Tel: 028 9031 1611. Textphone/Text Relay: 18001 028 9031 1611. www.publichealth.hscni.net 08/10